



OPEN ACCESS

EDITED BY

Pedro Gutierrez-Castrellon,
International Scientific Council for Probiotics
A.C., Mexico

REVIEWED BY

Maria Oana Sasaran,
"George Emil Palade" University of Medicine,
Pharmacy, Sciences and Technology of Târgu
Mureș, Romania
Amosy E. M'Koma,
Meharry Medical College,
United States

*CORRESPONDENCE

Batya Weiss

✉ batya.vais@sheba.health.gov.il

Nurit Loberman Nachum

✉ nurit.nachum@sheba.health.gov.il

RECEIVED 22 November 2023

ACCEPTED 12 September 2024

PUBLISHED 27 September 2024

CITATION

Nachum NL, Salomon N, Yerushalmy-Feler A,
Weintraub Y, Yogev D, Granot M, Haberman Y,
Ben-Horin S and Weiss B (2024) The efficacy
of curcumin/Qing Dai combination in children
with active ulcerative colitis: a multicenter
retrospective cohort study.
Front. Pediatr. 12:1342656.
doi: 10.3389/fped.2024.1342656

COPYRIGHT

© 2024 Nachum, Salomon, Yerushalmy-Feler,
Weintraub, Yogev, Granot, Haberman, Ben-
Horin and Weiss. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

The efficacy of curcumin/ Qing Dai combination in children with active ulcerative colitis: a multicenter retrospective cohort study

Nurit Loberman Nachum^{1,2*}, Nir Salomon³,
Anat Yerushalmy-Feler^{2,4}, Yael Weintraub^{2,5}, Dotan Yogev^{6,7},
Maya Granot^{1,2}, Yael Haberman^{1,2,8}, Shomron Ben-Horin^{2,3} and
Batia Weiss^{1,2*}

¹Pediatric Gastroenterology and Nutrition Unit, Edmond and Lily Safra Children's Hospital, Tel Hashomer, Israel, ²Sackler Faculty of Medicine, Tel Aviv University, Tel-Aviv, Israel, ³Department of Gastroenterology, Sheba Medical Center, Ramat Gan, Israel, ⁴Pediatric Gastroenterology Institute, Dana-Dwek Children's Hospital, Tel Aviv Sourasky Medical Center, Tel Aviv, Israel, ⁵Institute of Gastroenterology, Nutrition and Liver Diseases, Schneider Children's Medical Center of Israel, Petach Tikva, Israel, ⁶Juliet Keidan Institute of Pediatric Gastroenterology and Nutrition, Shaare Zedek Medical Center, Jerusalem, Israel, ⁷The Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel, ⁸Department of Pediatrics, Cincinnati Children's Hospital Medical Center, University of Cincinnati College of Medicine, Cincinnati, OH, United States

Background: Curcumin and Qing Dai (QD) are herbal extracts that recently showed efficacy in treating inflammatory bowel disease (IBD). Since 2016, a combination of curcumin with QD (CurQD) has been employed in our center for management of active ulcerative colitis (UC).

Objectives: We report the effectiveness and safety of CurQD therapy in children with mild-moderate UC or IBD-unclassified (IBD-U).

Design: A multicenter retrospective study.

Methods: Children aged ≤OP18 years who were treated with CurQD during 2017–2021 were included. Disease activity measures were Pediatric UC Activity Index (PUCAI), and fecal calprotectin (FC). The primary outcome was a decrease in PUCAI by ≥10 points, FC normalization (≤100 µg/gr when baseline ≥300 µg/gr) or a ≥50% decrease in FC.

Results: Of 30 patients (60% males, mean age 14 ± 3.9 years), 15 (50%), 13 (43%), and 2 (7%) had pancolitis, left-sided colitis and proctitis, respectively. The daily medication dose was 0.5–3 gm QD with 1–4 gm curcumin. Concomitant treatment at induction was corticosteroids (19%), biologics (28%) and 5-aminosalicylic acid (40%). The mean duration of induction was 11.6 weeks [95% confidence interval (CI) 10.2–13.1, range 8–16]. PUCAI decreased from a mean of 31.3 (95% CI 26.6–36.0, range 5–60) to 10.9 (95% CI 7.6–14.4, range 5–35) ($n = 26$, $p < 0.001$). FC response and normalization occurred in 11/12 and 7/12 patients, respectively. The median decline in FC was from 749 µg/gm [interquartile range (IQR) 566–1000] to 39 µg/gm (IQR 12–132) ($n = 15$, $p = 0.04$). During follow-up (median 8 months, IQR 6–10), 10 patients (33%) flared; five of them regained remission or responded to a treatment change. Of 18 patients treated beyond induction, 12 (67%) achieved clinical response and 10 achieved clinical remission by the end of follow up.

Conclusion: CurQD may be effective and safe as an add-on option to conventional management, for induction and maintenance in children with mild-moderate UC/IBD-U.

KEYWORDS

Qing Dai, curcumin, ulcerative colitis, pediatric, management

Introduction

Ulcerative colitis (UC) is a chronic inflammatory disease of the colonic mucosa, extending from isolated rectal disease to pancolitis. While the armamentarium for management of active UC has increased, novel compounds that may provide effective and safe treatment should still be explored, especially for the pediatric population (1–4).

Qing Dai (QD), also known as Indigo Naturalis, is a traditional Chinese medicine, derived from dried leaves and stems of blue flowers such as *Baphicacanthus cusia* Bremek, *Polygonum tinctorium* Ait and *Isatis indigotica* Fortune (5). QD has been used in traditional Chinese medicine for many years, due to its hemostatic, antipyretic, anti-inflammatory, sedative, antibacterial, antiviral, antioxidant and anti-tumor activities (6). QD has been shown to reduce inflammation through activation of the aryl hydrocarbon receptor (AhR), which promotes up-regulation of IL-22 and IL-10 (7). QD has been described as efficacious for treating inflammatory diseases like psoriasis (5, 8) and UC (9, 10).

A number of investigators have described the efficacy of a gut-directed formulation of curcumin in inducing and maintaining remission in patients with mild to moderate active UC (11, 12). Since 2016, a combination of curcumin with QD (CurQD) has been employed in our center, for management of active UC (13, 14). A recently published randomized placebo-controlled trial showed the efficacy of this combination in adults with active UC (15). However, data on curcumin or QD treatment in pediatric UC are scarce. Therefore, we aimed to report the efficacy and safety of CurQD management in a cohort of pediatric patients with UC or inflammatory bowel disease unclassified (IBD-U) who failed previous management.

assessed by the PUCAI (16) and by fecal calprotectin (FC) (17). We excluded from the analysis, patients who received less than 2 weeks of CurQD treatment, patients with missing significant data, and patients with loss of follow up.

Management regimen

CurQD is produced by Evinature (Binyamina, Israel) in a Good Manufacturing Practice facility. This management was provided to all the patients by a trained therapist of Chinese medicine, at the Gastroenterology Department. The doses were in the range of 0.5–3 gm/day for QD and 1–4 gm/day for curcumin, depending on the severity of the disease activity and the patient's weight, and based on published literature. While gut-directed curcumin was demonstrated to be effective in mild-moderate active UC (11), Qing Dai was shown to be effective for moderate-severe UC (18). Therefore, varying the ratios between these two compounds enables personalizing CurQD administration to each patient's condition. For example, a patient presenting with mild-moderate disease activity received CurQD with a lower QD dose and a higher curcumin dose. In contrast, a patient with more active disease received a higher QD and lower curcumin dose. The dose range was adjusted to pediatrics by slightly lowering the dose described in the above-cited adult studies. The CurQD management regimen generally followed an induction dose of CurQD for 16 weeks, after which the dose of QD was gradually reduced, until only gut-directed curcumin was continued. For patients who flared during the tapering, the minimal dose of QD within the combination that was needed to control the disease was continued.

Materials and methods

Patients

We conducted a multicenter retrospective cohort study of children treated at four pediatric gastroenterology units: Sheba Medical Center, Dana Dwek Children's Hospital, Schneider Children's Medical Center and Shaare Zedek Medical Center. Patient inclusion criteria were age ≤ 18 years; a Pediatric UC Activity Index (PUCAI) indicating mild or active UC or IBD-U (in the range of 10–35 and 40–60, respectively); and management with CurQD during 2017–2021. All the patients received CurQD at Sheba Medical Center, but clinical and endoscopic follow-ups were performed at each center. Clinical, laboratory and endoscopic data were retrieved from the medical records. Disease activity was

Outcome

The primary outcome was a clinical response after CurQD induction, up to 16 weeks from the beginning of management. This response was defined as a decrease in PUCAI by at least 10 points, and reduction in FC levels by 50%. Endoscopic MAYO scores prior to and post treatment were compared when available. The secondary outcome was remission (PUCAI ≤ 10 ; FC ≤ 100 , considering a baseline ≥ 300). Treatment failure was considered as discontinuation of CurQD due to adverse events or lack of response, hospitalization or induction of a biologic treatment or corticosteroids. When available, data regarding therapy beyond week 16 and until the end of the follow up were extracted. Concurrent therapy was given at the discretion of the treating physician.

Statistical analysis

Nonparametric data are presented as means with standard deviations or confidence intervals, or as medians and interquartile ranges (IQR). A Mann-Whitney test was used to analyze differences between nonparametric data. Missing data were considered as non-available.

The study was approved by the ethics committees of all the institutions (No. 8998-21-SMC).

Results

Patients and management

The 30 patients included were with mild (PUCAI 10–35) or moderate (PUCAI 40–60) active UC, or IBD-U at recruitment. Eighteen (60%) were males. Eleven patients were from Sheba Medical Center, nine from Dana Dwek Children’s Hospital, six from Schneider Children’s Medical Center and four from Shaare-Zedek Medical Center. **Table 1** presents the demographic and clinical characteristics of the study group. The mean age at CurQD initiation was 14 ± 3.9 years, and the median disease duration until CurQD treatment was 1.5 years (IQR 1.0–2.6). Disease extent was pancolitis in 15 (50%), left-sided colitis in 13 (43%) and proctitis in two (7%) patients. Eighteen patients (60%) were naïve to biologic treatment, and seven (22%) received more than one biologic therapy. At the time of CurQD induction, 19 (63%) patients were treated with 5-aminosalicylic acid, of them 13 as a single treatment. Six (20%) patients were treated with corticosteroids and six (20%) with immunomodulators. Nine (30%) patients were with active disease while receiving biologic treatment (4 Infliximab, 4 Vedolizumab and 1 Golimumab), to which the herbal compound was then added. The daily QD dose within the CurQD ranged from 0.5 to 3 gm, according to disease severity, combined with 1–4 gm of gut-directed curcumin.

The efficacy of induction

PUCAI at baseline showed mild activity (≤ 35) in 18 (60%) and moderate activity (PUCAI 40–60) in 8 (27%) patients. The mean PUCAI score decreased from 31.3 (95% CI 26.6–36.0, range 5–60) at baseline, to 10.9 (95% CI 7.6–14.4, range 5–35), at the end of induction (**Figure 1A**). The mean duration of induction was 11.6 weeks (95% CI 10.2–13.1, range 8–16) ($p < 0.001$). After induction, PUCAI decreased in 21 patients, showed no change in four patients and increased in one patient (**Figure 1B**). *Of the 18 patients with mild disease, the PUCAI score decreased by >10 points in 13 (72%), was not changed in 4 (22%) and increased in 1 (6%). PUCAI decreased by >10 points in all eight patients with moderate disease and available data.* In four patients, PUCAI was available at baseline and not post induction.

Among the 12 patients with paired FC measurements before and after induction, the median value declined from 749 $\mu\text{g/gm}$

TABLE 1 Demographic and clinical characteristics of pediatric patients with ulcerative colitis.

Age at diagnosis (years)	
Mean (SD)	12.17 (3.73)
Median (IQR)	13.15 (11–14)
Age at CurQD initiation (years)	
Mean (SD)	14 (3.89)
Median (IQR)	14.4 (12–16.5)
Gender	
Male (%)	18 (60)
Diagnosis (%)	
UC	28 (93)
IBD-U	2 (7)
Disease location (%)	
Proctitis	2 (7)
Left sided	13 (43)
Pancolitis	15 (50)
Previous treatments	
5ASA	29
Corticosteroids	19
Immunomodulators	8
Infliximab	10
Adalimumab	1
Vedolizumab	6
Golimumab	2
Concomitant treatment	
5ASA	19
Corticosteroids	6
Immunomodulators	6
Infliximab	4
Vedolizumab	4
Golimumab	1
Biologic treatment (%)	
None	20 (66)
1 Biologic	5 (17)
>1 Biologic	5 (17)
Treatment duration (months)	
Mean (SD)	12.9 (10.9)
Median (IQR)	11 (6–14)

UC, ulcerative colitis; IBD-U, inflammatory bowel disease unclassified; 5ASA, 5-aminosalicylic acid; SD, standard deviation; IQR, interquartile range.

(IQR 566–1000) to 39 $\mu\text{g/gm}$ (IQR 12–132) ($p = 0.04$) (**Figure 2C**). Eleven (91%) of these patients had an FC response at the end of induction. For seven (64%) of them, FC at baseline was $\geq 300 \mu\text{g/gm}$ and reached normalization ($\leq 100 \mu\text{g/gm}$). Five patients had FC levels higher than 100 $\mu\text{g/gm}$. Two of them had a clinical flare with endoscopic evidence of inflammation, and one had only endoscopic evidence of inflammation without clinical signs. FC improved in six patients with mild disease and in three with moderate disease. No changes were observed in serum hemoglobin, albumin or C-reactive protein levels before and after induction (**Figure 2**).

Three (10%) patients discontinued CurQD treatment during induction due to lack of efficacy. One of them was treated for only 2 weeks, and two had clinical and endoscopic evidence of active disease and were treated with biologic therapy.




FIGURE 1

Clinical response to treatment. Changes in Pediatric Ulcerative Colitis Activity Index (PUCAI) scores after CurQD induction. (A) Decline in mean PUCAI from baseline. (B) Trends of PUCAI in the study group **** $p < 0.0001$; NA, not available.

Efficacy of maintenance

Twenty-seven patients continued to maintenance treatment; the mean follow-up time was 14.5 ± 11 months. Of the 21 who responded to induction, eight (33%) had a flare, after a mean of $12 (\pm 10.6)$ months. One of these had a spontaneous remission with no additional treatment, one started 5-aminosalicylic acid medication, and one started budesonide with a good response. One patient temporarily discontinued CurQD, received antibiotics and renewed CurQD therapy after regaining remission. Another patient refused treatment escalation.

Data of paired FC at the end of the maintenance follow up were available for 13 patients. Twelve of them showed FC response, and five achieved normalization of FC. The median FC decreased from 749 (IQR 573–1,000) to 121.5 (IQR 35–174).

Colonoscopy was performed in eight (27%) patients during the follow-up period. Three of them had normal examinations and five had evidence of inflammation (Mayo score 2–3). One of the five patients improved clinically after the colonoscopy, while continuing management with CurQD only (Table 2).

Based on the available data, during the follow up, 19/30 (63%) patients did not change their concomitant medications, five patients changed or added biologic treatment, and six patients stopped or added other medication.

Safety

The adverse events included headache in one patient, which did not require any changes in medication and resolved spontaneously; and hematuria in one patient, which resolved after discontinuation

of curcumin. During the maintenance treatment, one patient had elevated transaminases, up to three times the upper normal limit, while having a concomitant Covid-19 infection. Although it was not clear whether the increase in enzyme levels resulted from the viral disease or was a side effect of the treatment, CurQD was discontinued, and the enzymes improved.

No subgroup analyses were performed due to the small sizes of the groups.

Discussion

Our findings showed potential efficacy and safety of CurQD management in children with UC or IBD-U who did not achieve complete remission with conventional management. Clinical response was achieved in 70% of the patients after induction; PUCAI and FC decreased significantly (Figures 1, 2C). Twelve of the 13 patients with available data showed a decrease in FC at the end of follow-up. We did not observe changes after induction in serum hemoglobin, albumin and C-reactive protein levels. This is probably because those markers were within normal limits in most of the patients prior to treatment induction.

In addition to efficacy, CurQD management seemed to be safe. The adverse events included only one patient with a spontaneously resolving headache, and one patient with elevated liver enzymes that co-existed with Covid-19 infection. That patient stopped treatment, and the liver enzymes normalized. However, it is not clear whether the increased levels were a side effect of the medication or due to the viral disease. We did not observe adverse effects that were previously reported for QD, such as gastrointestinal symptoms, liver dysfunction, cutaneous symptoms,




FIGURE 2

Laboratory response to treatment. Laboratory tests at baseline and at the end of induction showed no significant changes in serum albumin (A), hemoglobin level (B) or C-reactive protein (CRP) (D); and a significant decrease in fecal calprotectin (C) **** $p < 0.0001$, ns, non-significant.

leukopenia, dizziness and reversible pulmonary hypertension (18–21). We also did not observe even mild adverse events, as were reported in a clinical trial of curcumin, including abdominal distension, nausea and a transient increase in the number of bowel movements (12). One of our patients had hematuria, which is not

among the side effects that have been reported for curcumin; the condition resolved after curcumin discontinuation.

The proposed mechanism of action of curcumin is reducing inflammation by downregulation of genes, such as PI3 K, related to oxidative stress and fibrogenesis pathways; and by decreasing

TABLE 2 Endoscopic findings pre and post CurQD treatment and changes in treatment changes.

Patient	Previous treatment	Treatment	Lower endoscopy before QD treatment	Treatment change	Lower endoscopy after induction	Treatment change	Last lower endoscopy under QD treatment	Clinical outcome
1	5ASA, Steroids, Azathioprine, Infliximab, Adalimumab, Vedolizumab	Golimumab		CurQD			left-sided Mayo-3	No response - treatment change to Ustekinumab
2	5ASA, Steroids, Azathioprine, Infliximab	Infliximab Azathioprine	rectum- Mayo 2-3, descending and transverse Mayo 2	CurQD	rectum- Mayo 3, descending and transverse Mayo 1	change to Vedolizumab	normal	remission
3	5ASA, Steroids, MTX, Infliximab	5ASA	30 cm of ulcerated and hyperemic mucosa, Mayo 2	CurQD			30 cm of erythema	response
4	5ASA, Steroids, Azathioprine, Infliximab, Vedolizumab	Vedolizumab		CurQD, stop Vedolizumab			active colitis Mayo 2-3 until transverse colon	clinical improvement with CurQD only
5	5ASA, Steroids, Infliximab	Infliximab	08/2019 sigmoidoscopy- colitis, Mayo 2-30 cm.	CurQD			normal	response
6	5ASA, Steroids, Infliximab, Vedolizumab	Vedolizumab5ASA		CurQD	severe pancolitis		severe pancolitis	No response
7	5ASA, Steroids, Azathioprine, Vedolizumab	5ASA, Azathioprine		CurQD			active proctitis	No response
8	5ASA, Steroids, Infliximab	None	Severe pancolitis	CurQD	severe left-sided colitis		active disease	No response

5ASA, 5-aminosalicylic acid, Cur, curcumin, MTX, methotrexate, QD, Qing Dai.

the activity of proinflammatory cytokines like IFN- γ , TNF- α , IL-1 and IL-8 (22). QD reduces inflammation by activating AhR, which promotes up-regulation of IL-22 and IL-10 (7).

QD has been used in traditional Chinese medicine for centuries, for various infectious and inflammatory conditions (18). Several studies have examined the efficacy of QD management in adults and children with UC. In one of the first prospective studies, Sugimoto et al. (10) demonstrated significant clinical and endoscopic responses, with a good safety profile, within 8 weeks of a 2 gm/day QD treatment in adults with moderate UC. Two years later, the INDIGO study group (18) performed a multicenter randomized placebo-controlled trial in adults with UC. They showed a clinical remission rate of 55% with a 1 gm/day dose, and 38% with a 2 gm/day dose after 8 weeks of treatment. These proportions were significantly higher than those of the placebo. The high remission rate was demonstrated even in patients with steroid dependence, or with previous biological or immunomodulatory treatment, similar to some of our patients.

The first large QD study in children with UC (21) was conducted in Japan and included more than 100 patients. Clinical remission was shown for more than 80% within 6 months of treatment; the relapse rate was 64% in patients who used QD for more than 6 months. Seven patients experienced adverse events, which is a much higher rate than observed in the current study. The events included pulmonary hypertension in one patient, enteritis in five patients and headache in one.

Our management protocol included both curcumin and QD for 16 weeks, and then tapering of QD until discontinuation, while curcumin was continued. The advantage of our protocol is the prevention of QD side effects (13). In addition, the combined treatment enabled administering lower doses of the two components, which act by different mechanisms and together provide a broader and stronger effect (15).

The study group reflects the pediatric UC population, characterized by active disease despite standard management. The multiple center design and prolonged follow-up period are strengths of the study. Nonetheless, the study has limitations stemming from the small sample size and the retrospective design, which limits the control over data completeness. Data were missing for some of the outcomes and the use of concurrent medications. In addition, endoscopic follow-up was performed predominantly in non-responding patients. This may have influenced the observed rate of mucosal healing, which contrasted with the findings of the adult study (15). We included patients with moderate disease who did not achieve complete clinical remission or who flared on corticosteroids or biological therapy. Those patients had mild-moderate active disease at the initiation of CurQD; however, their initial disease state was moderate as is evident also from the proportion that failed biologics. The remaining patients had a mild-moderate disease, which indicates heterogeneity of the patient population.

Our report of minimal adverse events suggests that CurQD has a good safety profile. However, due to the retrospective design, some side effects may not have been reported. In addition, as echocardiograms were not performed routinely, pulmonary

hypertension, a known adverse event, may have occurred but not been reported.

In conclusion, our study shows that CurQD might be an effective and safe management for pediatric patients with mild to moderate UC who failed conventional management. Larger prospective studies are needed to evaluate the clinical and endoscopic effects of CurQD in children.

Data availability statement

The data analyzed in this study is subject to the following licenses/restrictions: retrospective data. Requests to access these datasets should be directed to batya.vais@sheba.health.gov.il.

Ethics statement

The studies involving humans were approved by Ethics committee Sheba medical center. The studies were conducted in accordance with the local legislation and institutional requirements. The ethics committee/institutional review board waived the requirement of written informed consent for participation from the participants or the participants' legal guardians/next of kin because This was a retrospective study.

Author contributions

NN: Data curation, Writing – original draft. NS: Conceptualization, Data curation, Writing – review & editing. AY-F: Data curation, Writing – review & editing. YW: Data curation, Writing – review & editing. DY: Data curation, Writing – review & editing. MG: Data curation, Writing – review & editing. YH: Writing – review & editing. SB-H: Conceptualization, Data curation, Methodology, Writing – review

& editing. BW: Conceptualization, Data curation, Methodology, Writing – review & editing.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Conflict of interest

Shomron Ben-Horin has received advisory board and/or consulting fees from Abbvie, Takeda, Janssen, Celltrion, Pfizer, GSK, Ferring, Novartis, Roche, Gilead, NeoPharm, Predicta Med, Galmed, Medial Earlysign, and Eli Lilly; and research support from Abbvie, Takeda, Janssen, Celltrion, Pfizer, & Galmed. Nir Salomon received speaking fees from Takeda, Bara Herbs and ITS Medical. The Sheba Medical Center has filed intellectual property on the combination of curcumin and QD. EviNature is a spin-off company of Sheba Medical Center. Shomron Ben-Horin & Nir Salomon received consulting fees and holds equity in EviNature.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Schroeder KW, Tremaine WJ, Ilstrup DM. Coated oral 5-aminosalicylic acid therapy for mildly to moderately active ulcerative colitis. A randomized study. *N Engl J Med.* (1987) 317(26):1625–9. doi: 10.1056/NEJM198712243172603
- Ford AC, Bernstein CN, Khan KJ, Abreu MT, Marshall JK, Talley NJ, et al. Glucocorticosteroid therapy in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol.* (2011) 106(4):590–9. quiz 600. doi: 10.1038/ajg.2011.70
- Gisbert JP, Linares PM, McNicholl AG, Mate J, Gomollon F. Meta-analysis: the efficacy of azathioprine and mercaptopurine in ulcerative colitis. *Aliment Pharmacol Ther.* (2009) 30(2):126–37. doi: 10.1111/j.1365-2036.2009.04023.x
- Bhattacharya A, Osterman MT. Biologic therapy for ulcerative colitis. *Gastroenterol Clin North Am.* (2020) 49(4):717–29. doi: 10.1016/j.gtc.2020.08.002
- Lin YK, Chang YC, Hui RC, See LC, Chang CJ, Yang CH, et al. A Chinese herb, indigo naturalis, extracted in oil (Lindioil) used topically to treat psoriatic nails: a randomized clinical trial. *JAMA Dermatol.* (2015) 151(6):672–4. doi: 10.1001/jamadermatol.2014.5460
- Gaitanis G, Magiatis P, Velegraki A, Bassukas ID. A traditional Chinese remedy points to a natural skin habitat: indirubin (indigo naturalis) for psoriasis and the malassezia metabolome. *Br J Dermatol.* (2018) 179(3):800. doi: 10.1111/bjd.16807
- Kawai S, Iijima H, Shinzaki S, Hiyama S, Yamaguchi T, Araki M, et al. Indigo naturalis ameliorates murine dextran sodium sulfate-induced colitis via aryl hydrocarbon receptor activation. *J Gastroenterol.* (2017) 52(8):904–19. doi: 10.1007/s00535-016-1292-z
- Lin YK, Chang CJ, Chang YC, Wong WR, Chang SC, Pang JH. Clinical assessment of patients with recalcitrant psoriasis in a randomized, observer-blind, vehicle-controlled trial using indigo naturalis. *Arch Dermatol.* (2008) 144(11):1457–64. doi: 10.1001/archderm.144.11.1457
- Suzuki H, Kaneko T, Mizokami Y, Narasaka T, Endo S, Matsui H, et al. Therapeutic efficacy of the qing Dai in patients with intractable ulcerative colitis. *World J Gastroenterol.* (2013) 19(17):2718–22. doi: 10.3748/wjg.v19.i17.2718
- Sugimoto S, Naganuma M, Kiyohara H, Arai M, Ono K, Mori K, et al. Clinical efficacy and safety of oral Qing-Dai in patients with ulcerative colitis: a single-center open-label prospective study. *Digestion.* (2016) 93(3):193–201. doi: 10.1159/000444217
- Lang A, Salomon N, Wu JC, Kopylov U, Lahat A, Har-Noy O, et al. Curcumin in combination with mesalamine induces remission in patients with mild-to-moderate ulcerative colitis in a randomized controlled trial. *Clin Gastroenterol Hepatol.* (2015) 13(8):1444–9 e1. doi: 10.1016/j.cgh.2015.02.019
- Hanai H, Iida T, Takeuchi K, Watanabe F, Maruyama Y, Andoh A, et al. Curcumin maintenance therapy for ulcerative colitis: randomized, multicenter, double-blind, placebo-controlled trial. *Clin Gastroenterol Hepatol.* (2006) 4(12):1502–6. doi: 10.1016/j.cgh.2006.08.008

13. Yanai H, Salomon N, Lahat A, Ungar B, Eliakim R, Kriger-Sharabi O, et al. Real-world experience with curcumin-QingDai combination for patients with active ulcerative colitis: a retrospective multicentre cohort study. *Aliment Pharmacol Ther.* (2023) 58(2):175–81. doi: 10.1111/apt.17538
14. Ben-Horin S, Kopylov U, Salomon N. Curcumin-QingDai combination as treatment for moderate-severe ulcerative colitis. *Case Rep Gastroenterol.* (2022) 16(3):563–8. doi: 10.1159/000526646
15. Ben-Horin S, Salomon N, Karampekios G, Viazis N, Lahat A, Ungar B, et al. Curcumin-QingDai combination for patients with active ulcerative colitis: a randomized double-blinded placebo-controlled trial. *Clin Gastroenterol Hepatol.* (2024) 22(2):347–56. doi: 10.1016/j.cgh.2023.05.023
16. Turner D, Otley AR, Mack D, Hyams J, de Bruijne J, Uusoue K, et al. Development, validation, and evaluation of a pediatric ulcerative colitis activity index: a prospective multicenter study. *Gastroenterology.* (2007) 133(2):423–32. doi: 10.1053/j.gastro.2007.05.029
17. Jukic A, Bakiri L, Wagner EF, Tilg H, Adolph TE. Calprotectin: from biomarker to biological function. *Gut.* (2021) 70(10):1978–88. doi: 10.1136/gutjnl-2021-324855
18. Naganuma M, Sugimoto S, Mitsuyama K, Kobayashi T, Yoshimura N, Ohi H, et al. Efficacy of indigo naturalis in a multicenter randomized controlled trial of patients with ulcerative colitis. *Gastroenterology.* (2018) 154(4):935–47. doi: 10.1053/j.gastro.2017.11.024
19. Orihara Y, Asakura M, Hida N, Kawai M, Sato T, Nishimura K, et al. Effect of oral qing-Dai medication on pulmonary arterial pressure levels in patients with ulcerative colitis. *Circ J.* (2020) 84(8):1339–45. doi: 10.1253/circj.CJ-19-1112
20. Adachi S, Nakano Y, Kondo T. Risk of pulmonary arterial hypertension during qing-Dai use for the treatment of ulcerative colitis. *Circ J.* (2020) 84(8):1235–6. doi: 10.1253/circj.CJ-20-0621
21. Kudo T, Jimbo K, Shimizu H, Iwama I, Ishige T, Mizuochi T, et al. Qing-Dai for pediatric ulcerative colitis multicenter survey and systematic review. *Pediatr Int.* (2022) 64(1):e15113. doi: 10.1111/ped.15113
22. Karthikeyan A, Young KN, Moniruzzaman M, Beyene AM, Do K, Kalaiselvi S, et al. Curcumin and its modified formulations on inflammatory bowel disease (IBD): the story so far and future outlook. *Pharmaceutics.* (2021) 13(4). doi: 10.3390/pharmaceutics13040484