



OPEN ACCESS

EDITED BY

Marwan El Ghoch,
University of Modena and Reggio Emilia, Italy

REVIEWED BY

Rizaldy Taslim Pinzon,
Duta Wacana Christian University, Indonesia

*CORRESPONDENCE

Guizhong Huang
✉ 14700010@qq.com

RECEIVED 09 October 2024

ACCEPTED 18 November 2024

PUBLISHED 04 December 2024

CITATION

Li X, Huang P and Huang G (2024)
Commentary: The roles of serum vitamin D
and tobacco smoke exposure in insomnia: a
cross-sectional study of adults in the
United States. *Front. Nutr.* 11:1500041.
doi: 10.3389/fnut.2024.1500041

COPYRIGHT

© 2024 Li, Huang and Huang. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Commentary: The roles of serum vitamin D and tobacco smoke exposure in insomnia: a cross-sectional study of adults in the United States

Xiangming Li, Peixing Huang and Guizhong Huang*

Shantou Hospital of Traditional Chinese Medicine, Guangzhou University of Traditional Chinese Medicine, Shantou, Guangdong, China

KEYWORDS

NHANES, data, question, vitamin D, tobacco smoke exposure

A Commentary on

The roles of serum vitamin D and tobacco smoke exposure in insomnia: a cross-sectional study of adults in the United States

by Gao, T., Hou, M., Wang, Q., Liu, D., Chen, F., Xing, Y., and Mei, J. (2023). *Front. Nutr.* 10:1285494. doi: 10.3389/fnut.2023.1285494

We read with great interest the article titled “The roles of serum vitamin D and tobacco smoke exposure in insomnia: a cross-sectional study of adults in the United States” by Gao et al. (1). The authors utilized data from the NHANES database and highlighted the potential regulatory role of serum vitamin D (VD) levels in the association between tobacco smoke exposure and insomnia. Their work offers valuable insights for the early identification and prevention of insomnia in high-risk populations. We commend the authors for shedding light on this important topic, which has potential implications for health policy.

However, we would like to raise a specific concern regarding the data used in the study. The authors mentioned that they utilized the NHANES data from 2005 to 2008. Upon review, we found that the data on serum 25(OH)D levels from the 2007 to 2008 NHANES cycle are currently unavailable for download (as stated on the NHANES website: <https://wwwn.cdc.gov/nchs/nhanes/continuousnhanes/default.aspx?BeginYear=2007>). We kindly request clarification on how the authors were able to access these data.

In addition, while we acknowledge the authors' comprehensive approach in controlling for various covariates (e.g., smoking, depression, OSA, work status, work shifts, and VD collection season), we would like to highlight some limitations of the study. As the study was based on a cross-sectional design, it could not establish causal relationships between VD levels, cotinine, and insomnia. Additionally, it could not conclusively determine the regulatory role of VD in the association between cotinine and insomnia.

We also wish to highlight that the study's reliance on self-reported insomnia might have introduced recall bias. Although clinical guidelines suggest that insomnia is often diagnosed through patient history rather than polysomnography (PSG), self-reported data can be subject to inaccuracies. In addition, the dietary intake information collected via two 24-h recalls might have further contributed to recall bias.

Furthermore, we would like to expand on the geographical limitations of the study. The findings from a U.S.-based cohort may not be directly applicable to regions with different seasonal and geographical characteristics. For instance, in tropical areas with only two distinct seasons, factors such as sunlight exposure, which affect vitamin D synthesis, may differ significantly from those in temperate climates. We recommend that the authors discuss this limitation and caution against extrapolating these findings to populations in regions with different climates and seasonal variations.

Moreover, the U.S. population sample used in this study had specific dietary patterns that may not align with those of populations in other countries. As dietary habits can significantly influence vitamin D levels, the conclusions drawn from this study may not be generalizable to regions with different dietary patterns. We encourage the authors to address this limitation and suggest future studies exploring these relationships in populations with varying dietary intake patterns.

We emphasize the need for additional prospective studies to better elucidate the causal relationship between VD and insomnia, as well as the potential role of VD in modulating the effects of tobacco smoke exposure on sleep disturbances.

Once again, we appreciate the authors' valuable contributions and look forward to seeing further research on this topic.

Author contributions

XL: Writing – original draft, Writing – review & editing. GH: Writing – review & editing. PH: Writing – review & editing.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Gao T. The roles of serum vitamin D and tobacco smoke exposure in insomnia: a cross-sectional study of adults in the United States. *Front Nutr.* (2023) 10:1285494. doi: 10.3389/fnut.2023.1285494